

What To Expect: Post Implant Placement Surgery

What To Do After Surgery?

Here are some guidelines and instructions for you to follow to aid in your healing and recovery:

1. Rest at home and take it easy for the first 24 hours following surgery.
2. You may experience light-headedness, dizziness, and sleepiness following surgery.
3. Please refrain from pulling on your lips or cheeks to inspect the surgical sites. This action could tear your sutures or delay healing.
4. Take your prescribed medications as directed.

How Much Swelling Is To Be Expected?

Significant swelling may occur after surgery. After the procedure, apply an ice pack on both sides of your face at 15 minute intervals for the remainder of the day. Moist heat such as a warm, wet wash cloth, will aid in reducing swelling after the first 48-72 hours.

Swelling usually reaches its maximum within the first 72 hours post-surgery and then slowly decreases.

Will There Be Bruising?

Bruising sometimes occurs after implant surgery. Bruising often develops a few days post-surgery and may be located either above or below the area that implants were placed. This is normal and typically resolves after the first week.

What Can I Eat After Surgery?

It is extremely important to be on a soft diet for the first 14 days of healing. During this time your implants(s) are integrating. Chew on opposite side if available. **AVOID:** Seeds, nuts, popcorn and hard foods

What Are Some Examples Of Acceptable Foods To Eat?

Acceptable foods are soft and easy to chew and swallow. These foods may be chopped, ground, mashed or pureed. This diet should be followed for the first 14 days after your surgery.

- **Fruits/Vegetables:** *Apple Sauce, banana, Cooked fruits or ripe fruits, Canned fruits without skin or seeds, Well-cooked soft veggies, or Mashed Potatoes*
- **Breads/Pasta/Rice:** *Pancakes, Macaroni/Pasta/Rice Pasta, Oat Meal/Cooked Cereal/Cream of Wheat, Waffles, Muffins*
- **Meats And Other Protein Sources:** *Broiled Fish, Roasted/Stewed Chicken, Soups with small soft pieces of vegetables and meat, Scrambled eggs ,and/or Tofu*
- **Other:** *Pudding/ Jell-O/ Yogurt, Ice Cream, Protein Shakes*

•

- **PLEASE:** Do not smoke for the first week after surgery. Smoking may dry out the tissues and delay healing. Smoking can also cause you to bleed more and for longer periods of time.
- **NOTE:** You can expect to be uncomfortable for the first 3 days after surgery. After the third day, discomfort generally lessens, with progressive improvement thereafter.
- **REMEMBER:** We are concerned about your well-being. If you should experience any difficulty in breathing, bleeding that you feel is excessive, persistent nausea or vomiting, or any pain that is not managed with prescribed medications, please contact us immediately. There is always a

doctor on call to answer your questions or to provide you with emergency care. 702-869-5700 – AFTER HOURS LINE PROVIDED.

How Do I Maintain Personal Hygiene?

You need to maintain a strong hygiene routine such as brushing, flossing and rinsing in order to help clean around your implant. After two weeks you may begin to use a water pick and floss.

Start antibiotics 2 days prior to surgery date- Amoxicillin 500mg

Take 1 tablet Motrin 800 mg, 1 hour prior to surgery