

What To Expect: Post All-On-4 Implant Placement Surgery

INFORMATION AND INSTRUCTIONS FOR THE ALL-ON-4® PROCEDURE

What To Do After Surgery?

Here are some guidelines and instructions for you to follow to aid in your healing and recovery:

1. Rest at home and take it easy for the first 72 hours following surgery.
2. You may experience light-headedness, dizziness, and sleepiness following surgery. A responsible adult should stay with you for 12 hours after surgery.
3. Please refrain from pulling on your lips or cheeks to inspect the surgical sites. This action could tear your sutures or delay healing.
4. Take your prescribed medications as directed.

How Much Swelling Is To Be Expected?

Significant swelling may occur after surgery. After the procedure, apply an ice pack on both sides of your face at 15 minute intervals for the remainder of the day. Moist heat such as a warm, wet wash cloth, will aid in reducing swelling after the first 48-72 hours.

Swelling usually reaches its maximum within the first 72 hours post-surgery and then slowly decreases.

Will There Be Bruising?

Bruising sometimes occurs after implant surgery. Bruising often develops a few days post-surgery and may be located either above or below the area that implants were placed. This is normal and typically resolves after the first week.

What Can I Eat After Surgery?

It is extremely important to be on a soft diet for the first 3 months of healing. During this time your implants are integrating, and you're wearing a temporary prosthesis that is not meant to be used to chew hard or crunchy foods.

What Are Some Examples Of Acceptable Foods To Eat?

Acceptable foods are soft and easy to chew and swallow. These foods may be chopped, ground, mashed or pureed. This diet should be followed for the first 3 months after your surgery.

Fruits/Vegetables:

- Apple Sauce
- Cooked fruits or ripe fruits
- Canned fruits without skin or seeds

- Well-cooked soft veggies

Meats And Other Protein Sources:

- Broiled Fish
- Roasted/Stewed Chicken
- Soups with small soft pieces of vegetables and meat
- Tofu

Breads/Pasta/Rice:

- Soft Loaf Bread
- Pancakes
- Muffins
- Waffles
- Macaroni/Pasta/Rice Pasta
- Oat Meal/Cooked Cereal/Cream of Wheat
- Mashed Potatoes

Other:

- Pudding/ Jell-O/ Yogurt
- Ice Cream
- Protein Shakes
- Scrambled eggs

How Do I Maintain Personal Hygiene?

An All-On-4® prosthesis sits firmly on your gums. Therefore, you need to maintain a strong hygiene routine such as brushing, flossing and rinsing in order to help clean under your prosthesis.

During the first week, use the provided monoject syringe to gently clean underneath your prosthesis. After two weeks you may begin to use a water pick and floss.

When Can I Return to My Daily Exercise Routine?

Refrain from exercise, heavy lifting and any physical activity for the first week following surgery. Elevation of heart rate and blood pressure may promote bleeding and delay healing. After the first week, and based on the doctor's discretion, patients may ease back in to physical activity.

PLEASE: Do not smoke for the first week after surgery. Smoking may dry out the tissues and delay healing. Smoking can also cause you to bleed more and for longer periods of time.

NOTE: You can expect to be uncomfortable for the first 3 days after surgery. After the third day, discomfort generally lessens, with progressive improvement thereafter.

REMEMBER: We are concerned about your well-being. If you should experience any difficulty in breathing, bleeding that you feel is excessive, persistent nausea or vomiting, or any pain that is not managed with prescribed medications, please contact us immediately. There is always a doctor on call to answer your questions or to provide you with emergency care. 702-869-5700 – AFTER HOURS LINE PROVIDED.

Start antibiotics 2 days prior to surgery date- Amoxicillin 500mg

Take 1 tablet Motrin 800 mg, 1 hour prior to surgery

Take 1 tablet Valium 5mg, 30 min prior to arrival at office – bring remaining to office..

Eat breakfast! Dress comfortable, stock up on soft foods, canned/bottled-meal supplements like ENSURE, SLIMFAST, or a similar substitute. So even if you don't feel like eating you will get proper nutrients those first few days.